

Sterile gel compositions for wound treatment**Publication number:** TW443932 (B)**Publication date:** 2001-07-01**Inventor(s):** HARDY CRAIG JULIAN [GB]; FINDLAY CHARLOTTE MARIA [GB]**Applicant(s):** JOHNSON & AMP JOHNSON MEDICAL [US]**Classification:**

- international: **A61K9/107; A61F13/00; A61K9/70; A61L2/07; A61L15/28; A61L15/44; A61L26/00; A61K9/107; A61F13/00; A61K9/70; A61L2/04; A61L15/16; A61L26/00; (IPC1-7): A61K47/10; A61K47/30**

- European: **A61L2/07; A61L15/28; A61L26/00B5; A61L26/00H2; A61L26/00H4; A61L26/00H7**

Application number: TW19950108358 19950811**Priority number(s):** GB19940014454 19940718**Also published as:**

 EP0693292 (A1)
 EP0693292 (B1)
 ZA9505607 (A)
 US5693624 (A)
 NO952824 (A)
 JP8187280 (A)
 GB2291348 (A)
 FI953453 (A)
 DE69524152 (T2)
 CA2153947 (A1)
 BR9503349 (A)
 AU2492895 (A)
 AU696405 (B2)
 AT209510 (T)

<< less

Abstract of TW 443932 (B)

The invention provides aqueous gel compositions for use as wound dressings and comprising from 2% to 10% w/v of alginate and from 25% to 40% w/v of polyhydric alcohol, said compositions being substantially sterile. The alginate is preferably sodium alginate and/or potassium alginate. The polyhydric alcohol is preferably propylene glycol or hexylene glycol. Also provided is a process of making such gel compositions comprising the steps of: providing an aqueous gel having the above composition and sterilising the aqueous gel. Preferably, the aqueous gel is sterilised by heating.

Data supplied from the esp@cenet database — Worldwide